

Statistical Analysis

Requested by Behavioral Health Program Manager, US Army.

[ADER: Abreaction, Desensitization and Emotional Reframing SM](#)
PTSD Checklist (PCL) Weathers et al., 1993

	Before	After	Time (Hours)	Calendar Days
Client 1	69	40	2	1
Client 2	63	42	12	16
Client 3	65	40	9	30
Client 4	65	40	0	1
Client 5	61	26	1	1
Client 6	65	34	4	14
Client 7	73	40	6	
Client 8	67	30	2	
Client 9	37	11	3	1
Client 10	63	33	4	10
Client 11	50	29	4	10
Client 12	45	29	6	
Client 13	65	30	2	1
Client 14	68	28	5	24
Client 15	70	31	6	16
Client 16	70	28	6	27
Client 17	70	26	4	16
Client 18	76	31	3	1
Client 19	75	35	4	14
Totals	1217	603	83	183
Average	64	32	5	10
	16.72	Two tailed T-test		

Greatest Impact: 2 hour sessions and multiple sessions per week appear to be most beneficial and time efficient. Values "post hoc" unless noted
Case Study report is available for each client.
<http://www.ptsd.va.gov/professional/pages/assessments/ptsd-checklist.asp>

TIME IS IMPORTANT

Trance State Therapy: In deepest relaxation, addressing Initial Sensitizing Events, building ego to compel forgiveness and resolution.

Management and Administration of the Abreaction:

Critical Care Intervention: Based on a document written by Paul Rieker with Roger A. Russell PhD

<http://www.BlessYourThoughts.com/ComplicatedGrief.doc>

- Objective: Compel the individual to "feel safe" in past events
- Behavior failures are believed to be based on Multiple Compounded Emotional Fixations through associational learning/sorting of emotions as response to sensitizing events, rather than "Systemic Breakdown", genetic failure or personality disorder.
- Trance Definition: elimination of peripheral distractions to achieve a single thought at a time.
- Trance state is achieved through management of respiration with other techniques.
- Sessions: accelerated and compacted
- Session includes sleep-time Emotional ReframingSM, applied in parallel.
- Therapy may be required due to, and after, years of emotional sorting/associational learning.
- Session best not limited to a simple schedule of 1 session per 1 hour per one week. Session may take as long as necessary to address the fixations, which become available during session.
- "Relapse" is misinterpreted. The reminder of an event a fixation, *which has not yet been addressed*, can trigger behavior failure.
- Potentially a permanent healing, unless of course, the individual could possibly become re-traumatized.
- Creates new and more brilliant emotional assignment to past hurt.

Bring the Study: We wish to further demonstrate research data for ADER (Abreaction, Desensitization and Emotional ReframingSM) to Post Traumatic Stress Disorder (PTSD) clients. Access to potential participants for the study, as well as a site for providing treatment and recording pre and post treatment outcomes. Please contact us using the information provided below.

Abreaction Desensitization and Emotional ReframingSM

SAFE CHILD: Leading the individual to Feel Safe in the memories contributing behavior of Post Traumatic Stress, Suicide, Complicated Grief and Depression.

5 Hour therapy program: Breaking as many as 40 emotional fixations, the possible change of behavior in the shortest period of time.

Treatment Research Opportunity

Paul Rieker Clinical Hypnotherapist (Practice beginning 1985)
Southern California 951-970-5641

www.BlessYourThoughts.com
BlessYourThoughts@gmail.com

Copyright 2011 Paul Rieker